


GOPHER FROG

Rana capito (formerly *R. areolata*)

Order:	Anura
Family:	Ranidae
FNAI Ranks:	G3G4/S3
U.S. Status:	None
FL Status:	Species of Special Concern



© Dan Hipes

Description: A medium-sized, boldly spotted frog with a chunky appearance: body short and plump, head large with somewhat rounded snout, legs relatively short. Back with somewhat warty skin and prominent, often bronze-colored longitudinal ridge on each side behind eye. Dorsal pattern of irregularly shaped dark spots on background that may be cream, gray, or brown. Chin and throat spotted, belly usually unmarked posteriorly. Adults 2.5 - 4 in. (63 - 102 mm) (excluding legs). Call resembles a deep snore. Tadpole large, to 3.5 in. (89 mm), globose, olive green, with large black spots on sides of tail.

Similar Species: Leopard frog (*Rana sphenocephala*), which may share breeding ponds with gopher frog, has large, dark brown spots on a green to

GOPHER FROG

Rana capito

brown background; however, body is more slender, snout very pointed, and throat and chin plain white. Tadpoles of the two species are very similar. Southern toad (*Bufo terrestris*) has dry, very warty skin, no raised ridges along edges of back, a pair of large raised glands behind eyes, and blunt snout. Spadefoot toad (*Scaphiopus holbrookii*) has vertical black pupils in golden eyes, dry skin, and a pair of hourglass-like lines rather than spots on back. All treefrogs have enlarged pads on toes.

Habitat: Dry, sandy uplands, chiefly sandhill and scrub, that include isolated wetlands or large ponds within about 1 mi. (1.7 km). Occasional in dry pine flatwoods, xeric hammock, and disturbed examples of above. Breeds chiefly in seasonally flooded, temporary ponds, but also in some permanent waters. Nocturnal, normally spending daytime in stumpholes, tunnels, or burrows, especially those of gopher tortoise (*Gopherus polyphemus*).

Seasonal Occurrence: Migrates to ponds for breeding from October through April, though may also breed during summer in central and southern Florida.

Florida Distribution: Most of state excluding Everglades and Keys; potential but not documented for some counties indicated on map. Two subspecies: dusky gopher frog (*R. c. sevosa*) in western panhandle, Florida gopher frog (*R. c. aesopus*) in peninsula and eastern panhandle.

Range-wide Distribution: Southeastern Gulf and Atlantic Coastal Plains, from North Carolina to eastern Louisiana.

Conservation Status: Many protected conservation lands in Florida support gopher frogs, although attention to managing and protecting breeding habitat and migratory pathways is often insufficient.

Protection and Management: Maintain large tracts of native vegetation in sandy, upland habitats that include wetlands. Allow fires to burn through dry wetland basins in addition to uplands. Manage uplands for gopher tortoises. See recommendations for striped newt (*Notophthalmus perstriatus*).

Selected References: Ashton and Ashton 1988a, Bartlett and Bartlett 1999, Conant and Collins 1991, Franz 1986, Franz and Smith 1999, Moler (ed.) 1992, Mount 1975.