

## CELESTIAL LILY

*Nemastylis floridana* Small


**Synonyms:** none

**Family:** Iridaceae (iris)

**FNAI Ranks:** G2/S2

**Legal Status:** US—Mgmt Concern FL—Endangered

**Wetland Status:** US—OBL FL—FACW



Alfred R. Schotz

**Field Description:** Perennial **herb** from a bulb with a single, tall, slender **stem**, occasionally branching on robust plants. **Basal leaves** few, grass-like, sometimes more than 2 feet long. **Stem leaves** small and scattered along the stem. **Flowers** more than 1.5 inches across, with 6 dark blue, spreading **petals and sepals (tepals)**; flowers open around 4 pm and close by dusk. **Stamens** with 3 coiled, yellow anthers; **style** divided into 6 narrow, pointed branches. **Fruit** an erect, oval capsule, about 0.5 inch long.

**Similar Species:** Blue-eyed grasses (*Sisyrinchium* spp.) are also in the iris family; they have much smaller blue flowers that are open throughout the day in the spring and summer. Celestial lily is the only iris-like species in FL to open in the late afternoon in the fall.

**Related Rare Species:** See Bartram's ixia (*Calydorea coelestina*), state-endangered, in this guide.

## Celestial lily

## *Nemastylis floridana*

**Habitat:** Wet flatwoods, prairies, marshes, cabbage palm hammocks edges.

**Best Survey Season:** Flowers from 4-6 pm, August–October.

**Range-wide Distribution:** Endemic to eastern counties of FL, primarily in the St. Johns River drainage.

**Conservation Status:** Once widespread in eastern FL, this species now occurs in about 15 managed areas, where it may be locally abundant if its habitat is frequently burned.

**Protection & Management:** Burn flatwoods and prairies every 2 - 3 years. Protect wetlands from draining, ditching, and conversion to pasture and pine plantation.

**References:** Coile 2000, Goldblatt 1975, Kral 1983, MacKiernan and Norman 1979, Small 1931b, Ward 1979, Wunderlin 1998, Wunderlin and Hansen 2000a.

