MODEST SPLEENWORT

Asplenium verecundum Chapman ex. L.

Underwood

Synonyms: Asplenium myriophyllum (Sw.) C.

Presl

Family: Aspleniaceae (spleenwort)

FNAI Ranks: G1/S1

Legal Status: US-none FL-Endangered





Field Description: Fern with tufted, lacy, evergreen **fronds**, usually drooping or arching, to 16 inches long, with a short, brownish-black **leaf stalk**. **Leaflets** alternate, in 8 - 22 well spaced pairs, leaflets divided into 2 - 5 segments, each with one **sorus** covered by an **indusium**.

Similar Species: Eared spleenwort (*Asplenium auritum*, drawing, lower right), state-endangered, has sharply toothed leaflets with an "ear" at the base of most leaflets and up to 10 pairs of sori per leaflet. Modest spleenwort hybridizes with a common species, cutleaf spleenwort (*Asplenium abscissum*), to produce a rare hybrid, Curtiss' spleenwort (photo, above left, *Asplenium x curtissii*). Cutleaf spleenwort backcrosses with Curtiss' spleenwort to produce another rare hybrid, ruffled spleenwort (see photo, above right, *Asplenium x plenum*). Also see American toothed spleenwort (*A. dentatum*) and single-sorus spleenwort (*A. monanthes*) in this guide.

Related Rare Species: None

Habitat: Rockland hammocks, limestone outcrops, grottoes, and sinkholes.

Best Survey Season: All year.

Range-wide Distribution: Modest spleenwort: endemic to FL. Eared spleenwort: FL, Mexico, West Indies, Central and South America.

Conservation Status: Most populations are on private lands where clearing, logging, and draining have destroyed the hammocks, swamps, and outcrops.

Protection and Management: Preserve swamp and hammock forests; protect and restore natural hydrology. Eradicate exotic pest plants. Enforce plant protection laws and prosecute plant poachers.

References: Coile 2000, FNA 1993, IRC 1999, Lellinger and Evans 1985, Nauman 1986, Nelson 2000, Ward 1979, Wunderlin 1998, Wunderlin and Hansen 2000a, Wunderlin and Hansen 2000b.