



E.B. glade (Gadsden County) with black bogrush in foreground

Photo by Ann F. Johnson

Upland Glade

Description: Upland glade is a largely herbaceous community with woody inclusions that occurs on thin soils over limestone outcrops on steep topography in Jackson and Gadsden counties. It is found in small openings ranging from 0.1 to 2 acres in size within an otherwise forested landscape. Open portions of upland glade are dominated by black bogrush (*Schoenus nigricans*) and/or other graminoids, such as poverty dropseed (*Sporobolus vaginiflorus*), hairawn muhly (*Muhlenbergia capillaris*), the stoloniferous form of little bluestem (*Schizachyrium scoparium*), pineywoods dropseed (*Sporobolus junceus*), and spreading beaksedge (*Rhynchospora divergens*), with variable amounts of limestone rock exposed. Other characteristic herbs include Cherokee sedge (*Carex cherokeensis*), lanceleaf tickseed (*Coreopsis lanceolata*), cypress witchgrass (*Dichanthelium dichotomum*), false garlic (*Nothoscordum bivalve*), Boykin's milkwort (*Polygala boykinii*), and starrush whitetop (*Rhynchospora colorata*). Areas of thinner soil or bare rock support diamondflowers (*Stenaria nigricans*), Gulf spike-moss (*Selaginella apoda* var. *ludoviciana*), a short turf-like moss (*Weissia jamaicense*), and blue-green algae (*Nostoc* sp.). A set of limestone-loving shrubs and trees on deeper soil within the glade, or on the edges form a shrubby transition to upland hardwood forest. Characteristic woody species include red cedar (*Juniperus virginiana*), eastern redbud (*Cercis canadensis*), sugarberry (*Celtis laevigata*), roughleaf dogwood (*Cornus asperifolia*), white ash (*Fraxinus americana*), yaupon (*Ilex vomitoria*), and winged elm

(*Ulmus alata*). Rattan vine (*Berchemia scandens*) is commonly found scrambling over these shrubs and small trees.

Upland glade was first mapped in the course of a survey of the Apalachicola ravines in Gadsden County (Leonard and Baker 1982). More glades have recently been mapped in Jackson County. In Gadsden County upland glade occurs on limestone outcrops of the Miocene Chattahoochee Formation (Rupert 1990; Means, pers. comm. 2007) primarily between the 90 and 120 foot contour lines. In Jackson County upland glade is found primarily on Oligocene Marianna Limestone (Moore 1955; Means, pers. comm. 2007) between the 130 and 150 foot contours.

Upland glade usually occupies areas too small to be distinguished at the scale of county soil survey maps. They may occasionally be shown as rock outcrop symbols within the matrix of the surrounding forest soils. In Gadsden County these matrix soils are Binnsville soils or Cuthbert, Boswell, and Susquehanna soils on moderate to steep slopes (Thomas et al. 1961). In Jackson County the matrix soils are the Oktibbeha variant rock outcrop complex (Duffee et al. 1979).

Characteristic Set of Species: black bogrush, poverty dropseed, diamondflowers, hairawn muhly, Boykin's polygala, red cedar

Rare Species: Ten northern species that are listed as rare in Florida, but range widely outside the state, occur on upland glades: green milkweed (*Asclepias viridiflora*), poppy mallow (*Callirhoe papaver*), small-toothed sedge (*Carex microdonta*), Carolina larkspur (*Delphinium carolinianum*), shootingstar (*Dodecatheon meadia*), eastern purple coneflower (*Echinacea purpurea*), little-people (*Lepuropetalon spathulatum*), pinnate-lobed coneflower (*Rudbeckia triloba* var. *pinnatiloba*), nettle-leaved sage (*Salvia urticifolia*), and shade betony (*Stachys crenata*). Eleven other species on upland glades that are rare in Florida but have widespread distributions outside the state also are being considered for listing by FNAI: meadow garlic (*Allium canadense* var. *mobile*), sideoats gramma (*Bouteloua curtipendula*), smooth oxeye (*Heliopsis helianthoides* var. *gracilis*), Small's ragwort (*Packera anonyma*), narrowleaf mountain-mint (*Pycnanthemum tenuifolium*), pinnate prairie coneflower (*Ratibida pinnata*), poverty dropseed (*Sporobolus vaginiflorus*), western silver aster (*Symphotrichum sericeum* var. *microphyllum*), Short's aster (*Symphotrichum shortii*), smooth blue aster (*S. laeve*), and hairyjoint meadowparsnip (*Thaspium barbinode*). Other possible candidates for listing include purple meadowparsnip (*Thaspium trifoliatum*) and golden alexanders (*Zizia aurea*).

Range: Upland glade is found only in Florida and adjacent Decatur County, Georgia (Ward and Gholson 1987) in two clusters about 30 miles apart. One cluster is in a four-square-mile area northwest of the town of Marianna in Jackson County, totaling 9 sites and 8.5 acres, and the other is an eighteen-square-mile area south and east of the town of Chattahoochee in Gadsden County, totaling 13 sites and 14.5 acres. New sites continue to be found within these areas.

Upland glade in Florida shares many species with other areas to the north and west where limestone is at or near the surface. These include the cedar glades on limestone outcrops in Tennessee, northern Alabama and northwest Georgia (Baskin and Baskin 1999), the

shallow-soil Black Belt prairies on Selma Chalk in Alabama and Mississippi (Schotz 2005), the Oaky Woods Wildlife Management Area in central Georgia (Echols, Jr. 2007), and various outcrop and prairie communities in Louisiana (MacRoberts and MacRoberts 1995; 1996) and southeastern Texas (Bridges et al. 1989). None of these areas, however, have black bogrush, a dominant species on Florida glades. In the United States this species is found only in Florida, Texas, California and Nevada; it also occurs in western Europe, South Africa, the West Indies and Mexico (Sparling 1968). The cedar glades of Tennessee and northern Alabama are dominated by the annual grass, poverty dropseed (Quarterman 1950; Baskin and Baskin 1999) and the Black Belt Prairies of Alabama and Mississippi by little bluestem (Schotz 2005). Both these species may also dominate portions of upland glade in Florida. Florida glades have the most species in common with the Black Belt Prairies in Alabama and the Oaky Woods prairies of Georgia which are the closest source of many calcium-loving species on Florida upland glades, being less than 100 miles distant. Species rare in Florida that are also found in these two areas include green milkweed, Carolina larkspur, and pinnate prairie coneflower; species shared with the Georgia Oaky Woods prairies also include smooth oxeye, Small's ragwort, and golden alexanders (Echols, Jr. 2007); those shared with the Black Belt Prairies in Alabama also include small-toothed sedge, nettle-leaved sage, and western silver aster (Schotz 2005)

Natural Processes: Soil depth is cited as the factor inhibiting tree and shrub colonization of herbaceous portions of limestone glades in Tennessee and Alabama (Quarterman 1950; Quarterman et al. 1993; Baskin and Baskin 1999). A small study measuring soil depth and pH on transects across three of the Gadsden County glades (Coultas 1983) lends support to this hypothesis for the Florida upland glades. Soil depth to hard limestone ranged from 4 to 14 inches in 13 samples taken along transects on the open herbaceous portions of three glades, with most samples falling in the 6 to 8 inch range. These figures are similar to soil depths measured on open glades in Tennessee (Baskin and Baskin 1999). Depth to limestone in four samples from the forest immediately surrounding the glades ranged from 21 to over 60 inches (Coultas 1983). Soils on the open glade were more alkaline (pH 7.5-7.6) than soil of the surrounding forest (pH 4.8-6.4; Coultas 1983). An additional stress on plants on glades is the alternation between wet and dry soils. Soils on the herbaceous portion of the glades are often wet and “mushy” in early spring and become nearly completely dry by fall.

Fire may have swept through upland glades at irregular intervals, especially those that were bordered by longleaf pine-dominated upland pine or sandhill communities that would have naturally tended to burn frequently.

Community Variations: Although upland glades in Jackson and Gadsden counties share dominant and many characteristic species, there is a set of characteristic species that serve to differentiate them. Present on Gadsden, but not Jackson County glades are poppy mallow, small-toothed sedge, littlehip hawthorn (*Crataegus spathulata*), orange coneflower (*Rudbeckia fulgida*), Gulf spike-moss, western silver aster, and Florida bully (*Sideroxylon reclinatum*). Present on Jackson but not Gadsden County glades are *Coreopsis* sp. (narrow, glabrous leaves), Carolina woollywhite (*Hymenopappus scabiosaeus*), Small's ragwort, pinnate-lobed coneflower, and gum bully (*Sideroxylon lanuginosum*). Two species are more common in one or the other county: Florida maple

(*Acer saccharum* var. *floridanum*) is more common around glades in Gadsden County; chinquapin oak (*Quercus muehlenbergii*) is more common around glades in Jackson County.

Associated Communities: The woody component of upland glade can be distinguished from the adjacent upland hardwood forest by the presence of red cedar, the concentration of other calciphilic species, and the short stature of the trees and shrubs. Upland glade can be distinguished from mesic flatwoods, which sometimes also has black bogruss in the ground layer where limestone is near the surface, by the presence of green milkweed and other characteristic northern calciphiles and the absence of saw palmetto (*Serenoa repens*), gallberry (*Ilex glabra*) and a pine canopy. Occasionally glade-like areas occur within the upland hardwood forest community where limestone is close to, but not at, the surface. These may be small grassy areas with a few of the characteristic glades herbs but without either black bogruss or the other short graminoid species as dominants.

Management Considerations: Although the ecology of the upland glade community has not been studied, it seems a reasonable working hypothesis that soil depth prevents woody colonization on the rockier portions of the glades, such as those areas dominated by diamond flowers and poverty dropseed, as has been proposed for areas of similar vegetation in the Tennessee glades (Quarterman 1950). Portions of the glades supporting taller grasses such as Gulf hairawn muhly and little bluestem may be more susceptible to woody encroachment, since increases in cover of red cedar have been observed at some glades over the course of several decades. Thus it may be that infrequent fires play a role in preventing woody species from shading out the characteristic graminoid dominants, particularly in those glades bordered by pine communities. Occasional droughts may also play a similar role in retarding woody growth on the glade. Casual observation of one burn on a glade indicated that the herbaceous species recovered to pre-burn levels of cover within a year or two, but the question of natural fire interval for this community needs more research. Growing season burns on experimental plots on a glade in Georgia also showed little effect on 14 herbaceous species of conservation concern two years later, but did result in a dramatic decline in populations of young trees on the glade (Duncan et al. 2008).

Threats to upland glades are primarily physical disturbance, since to non-botanists they appear indistinguishable from an artificial clearing in the forest. Only three of the twenty-two known glades are on public lands. Known glades have been lost to road widening, plowing for a game food plot, and limestone mining. In the past, others were probably destroyed by clearing, dumping, and disturbance from being used for parking logging machinery and as staging areas for timber operations. When the soil is disturbed, glades are vulnerable to being colonized by a dense sward of the exotic centipede grass (*Eremochloa ophiuroides*) which effectively precludes re-colonization by native glades species. Other exotics invading disturbed glades include Formosa firethorn (*Pyracantha koidzumii*), and Chinese and glossy privets (*Ligustrum sinense*, *L. lucidum*).

Exemplary Sites: The best developed glades are on private lands (Chalky, Brickyard, and E.B. glades in Gadsden County and Brooks 1 in Jackson County). Only a few glades are on public lands: one small glade in Florida Caverns State Park (Jackson County), and several small glades on lands of the Florida Department of Corrections south of US 90,

about one mile east of Chattahoochee; the latter are not specifically managed for protection of their natural resources.

Global and State Rank: G1/S1

Crosswalk and Synonyms:

SAF 73/Southern Redcedar

Other synonyms: cedar glades (Baskin and Baskin 1999), limestone glades (Bridges and Orzell 1986)

References:

Baskin, J.M., and C.C. Baskin. 1999. Cedar glades of the southeastern United States. Pages 206-219 in R.C. Anderson, J.S. Fralish, and J.M. Baskin, editors. *Savannas, Barrens, and Rock Outcrop Plant Communities of North America*. Cambridge University Press, Cambridge.

Bridges, E.L., and S.L. Orzell. 1986. Distribution patterns of the non-endemic flora of Middle Tennessee limestone glades. *Association of Southeastern Biologists Bulletin* 33:155-166.

Bridges, E.L., S.L. Orzell, and L.C. Anderson. 1989. *Carex microdonta* Torr. & Hook. (Cyperaceae) new to Florida. *Sida* 13:379-381.

Coultas, C.L. 1983. Examination of soils associated with four small "limestone glade communities" surrounded by forest in the hills of western Gadsden County. Florida Natural Areas Inventory, Tallahassee.

Duffee, E.M., W.J. Allen, and H.C. Ammons. 1979. Soil survey of Jackson County, Florida. United States Department of Agriculture, Soil Conservation Service in cooperation with the University of Florida Institute of Food and Agricultural Sciences, Agricultural Experiment Stations, Soil Science Department, Gainesville, Florida.

Duncan, R.S., C.B. Anderson, H.N. Sellers, and E.E. Robbins. 2008. The effect of fire reintroduction on endemic and rare plants of a Southeastern glade ecosystem. *Restoration Ecology* 16:39-49.

Echols, S.L., Jr. 2007. Vascular flora of remnant blackland prairies and associated vegetation of Georgia. Masters Thesis, University of Georgia, Athens.

Leonard, S.W., and W.W. Baker. 1982. Biological survey of the Apalachicola Ravines biotic region of Florida. Report to the Florida state office of The Nature Conservancy and to the Florida Natural Areas Inventory, Tallahassee, Florida.

MacRoberts, B.R., and M.H. MacRoberts. 1995. Vascular flora of two calcareous prairie remnants on the Kisatchie National Forest, Louisiana. *Phytologia* 78:18-27.

- MacRoberts, B.R., and M.H. MacRoberts. 1996. The floristics of calcareous prairies on the Kisatchie National Forest, Louisiana. *Phytologia* 81:35-43.
- Means, H. Geologist with the Florida Geological Survey, Tallahassee. Personal Communication. 2007
- Moore, W.E. 1955. Geology of Jackson County, Florida: Florida Geological Survey Bulletin 37:1-101.
- Quarterman, E. 1950. Major plant communities of Tennessee Cedar Glades. *Ecology* 31:234-254.
- Quarterman, E., M.P. Burbank, and D.J. Shure. 1993. Rock outcrop communities: limestone, sandstone, and granite. Pages 35-86 in W.H. Martin, S.G. Boyce, and A.C. Echternacht, editors. *Biodiversity of the Southeastern United States: Upland Terrestrial Communities*. John Wiley and Sons, Inc., New York.
- Rupert, F.R. 1990. Geology of Gadsden County, Florida. Florida Geological Survey, Tallahassee Florida.
- Schotz, A. 2005. A checklist of vascular flora in the Black Belt Prairies of Alabama. Alabama Natural Heritage Program, Montgomery, Alabama.
- Sparling, J.H. 1968. Biological flora of the British Isles: *Schoenus nigricans* L. (*Chaetospora nigricans* Kunth). *Journal of Ecology* 56:883-899.
- Thomas, B.P., H.H. Weeks, and M.W. Hazen, Jr. 1961. Soil Survey of Gadsden County, Florida. United States Department of Agriculture, Soil Conservation Service in cooperation with the University of Florida Institute of Food and Agricultural Sciences, Agricultural Experiment Stations, Soil Science Department, Gainesville, Florida.
- Ward, D.B., and A.K. Gholson. 1987. The hidden abundance of *Lepuropetalon spathulatum* (Saxifragaceae) and its first reported occurrence in Florida. *Castanea* 52:59-67.